

## Research Article

## KINETIC CHARACTERISTICS OF UREASE ISOLATED FROM SELECTED BEANS SPECIES AS AFFECTED BY GERMINATION

<sup>1</sup>Ogunmolasyi Adewoyin M. and \*<sup>1</sup>Evans C. Egwim

\*Department of Biochemistry, Federal University of Technology, Minna, Nigeria

<sup>1</sup>Department of Biochemistry, Federal University of Technology, Minna, Nigeria\*Corresponding: [evanschidi@gmail.com](mailto:evanschidi@gmail.com), Tel: +2347065809474Accepted 23<sup>rd</sup> December 2011

## ABSTRACT

In this research four different beans (*Glycine max*, *Mucuna pruriens*, *Kerstings' geocarpa* and *Vigna mungo*) were germinated in order to evaluate the implication of germination on the kinetic parameters ( $V_{max}$  and  $K_m$ ) of urease isolated from the various beans and compared to the non-germinated (control). The results showed that,  $V_{max}$  and  $K_m$  of urease from germinated beans increased significantly when compared ( $p < 0.05$ ) to urease from the control. This simply means that, urease from germinated beans demonstrated higher activity with lower substrate sensitivity when compared ( $p < 0.05$ ) to the control.

**Keywords:** Kinetics, Germination, Urease.

## INTRODUCTION

Urea is efficiently hydrolyzed by urease (E.C 3.5.15) a nickel dependent enzyme, to form ammonia and carbon (IV) oxide. The enzyme urease was first isolated and crystallized by James Summer in 1926 from jack beans (*canavalia ensiformis*) as a pure crystalline enzyme (James, 1926). Crystalline form of urease was the first obtained for a known enzyme.

Urea fertilizer is the predominant solid nitrogen (N) fertilizer in Chinese agriculture, and constitutes over 50% national nitrogenous fertilizer consumption (Zhu and Chen, 2002). However, the reaction catalyzed by urease is essential to make urea nitrogen available for plants growth and development (Gerenda's et al., 1999). This is as a result of the fact that urea is rich in nitrogen and urease gets the N available for plants.

Besides, urease is a very efficient way of determining urea level in biological fluid as well as in drinking and industrial water. Additionally, it is also used as reducing agent in alcoholic beverages (Fujinawa and Dela, 1990; Fujinawa and Ouch, 1991). Urease is also found useful since urea has commercial uses for the manufacture of hair conditioner, glues, plastic and animal feed as a browning agent in factory produced pretzels and also drug uses of urea in topical dermatological products, for nonsurgical debridement of nails and as a diuretic (Wikipedia, 2006). In the assessment of environmental pollution with mercury (II), array based

enzyme biosensors have been reported for screening various environmental pollutants, mainly heavy metals and pesticides. Urease has been extensively used as a model enzyme to elucidate the applicability of inhibitory assay for mercury (II) (Tsai and Doong, 2005; Preinmger, 1999 and Mahnaod, 2001).

This work was designed to investigate the kinetic characteristics of urease from germinated beans when compared to the non-germinated.

## Materials and Methods

Soya bean-*glycine max*, Black gram-*Vigna mungo*, *Mucuna* spp- *Mucuna pruriens* and *Kerstingella-Kerstings geocarpa* were procured from local market. Nessler's reagent, tris buffer and other chemical used were of analytical grade.

## Methods

## Germination Process

The beans were allowed to germinate or to develop the beginning of a root system. This was achieved in 3 days. Beans are essentially a seed and under warm and moist condition will begin to sprout or root, just as it would if it were growing in nature. Under this humid condition, bean seed initially swelled than germinated such that the radicle appeared. The germination process was stopped by freezing the beans for 12 hours followed by drying.

## Urease Isolation

Individual germinated and non-germinated beans were blended coarsely with laboratory mortar and pestle (wooden) and 50g of the same beans were soaked overnight in 100ml extraction buffer (0.025m Tris-acetate buffer, pH 6.5) at 4°C (Refrigeration temperature). The pH range for Tris buffer (7.5-9.5), is higher than the chosen pH 6.5; this was done for purification reason and to enhance maximum extractable activity and higher percentage recovery of enzyme during purification (Kayastha and Nilanjana, 1999). The next day, the soaked beans meals were swirled for 2 minutes and then sieved with four layers of pre-washed and dried white handkerchief, and the filtrate centrifuged

at 15000 rev/min for 15 mins at 4°C. The clear supernatant was collected while the sediment was discarded.

### Protein Determination

The protein content of the beans was determined spectrophotometrically after precipitation with acetone (Whitaker and Granum, 1980).

### Urease assay

To measure the urease activity, the ammonia liberated on incubating the enzyme with urea for 30 mins was determined using Nessler's reagent (Kulshrestha and Husain, 2006). One unit of urease activity liberates 1.0 mol of ammonia per min from 0.10 M urea under standard assay conditions.

The urease from both germinated and the control beans were characterized for the following parameter under standard condition of assay;

- pH.
- Temperature.
- Substrate concentration and enzyme kinetics ( $K_m$  and  $V_{max}$ ).

### pH Stability

In order to determine the effect of pH on activity of urease from germinated and control beans, pH values of 0.05M Tris-acetate buffer ranged between 5.5 and 8.0 and incubated at constant temperature 37°C and 0.2M urea.

### Thermal Stability

To determine the optimal temperature up to which the urease enzyme from both germinated and the control beans can withstand thermal stress, free urease enzyme was suspended in Tris-acetate buffer (0.05 M) and optimum pH

above, and incubated at different temperatures (30 to 80 °C) for 30 min before the activity was measured.

### Determination of the Kinetic Parameters

The kinetic parameters  $K_m$  and  $V_{max}$  were calculated by Lineweaver-Burk equation chosen according to the properties of substrate concentration, the linear transformation of Michaelis-Menten equation (Segel, 1975):


$$\frac{1}{V} = \frac{K_m}{V_{max}} \cdot \frac{1}{[S]} + \frac{1}{V_{max}}$$




Where  $V$  is the enzyme reaction velocity,  $[S]$  is the concentration of substrate (mmol L<sup>-1</sup>),  $K_m$  is the Michaelis constant (mmol L<sup>-1</sup>).  $V_{max}$  is the maximum enzyme reaction velocity ( $\mu\text{mol/s/mg protein}$ )  $K_m$  indicates the affinity of urease to its specific substrate urea, and gives the substrate concentration at which the reaction rate reaches half of its maximum value ( $V_{max}/2$ ). The rate of the reaction was measured as  $\mu\text{moles of ammonia produced} / \text{s} / \text{mg enzyme}$ .

### Result and Discussion

Experimental results about urease activity from both germinated and non-germinated were reported. The results showed that germination had significant influence on urease kinetic parameters when compared ( $P < 0.05$ ) to the control, under standard condition of assay (Table 2).

The table 1.0 below shows the result of optimum pH, temperature and substrate concentration for maximum hydrolysis of urea by urease isolated from germinated and control (non-germinated) of the various beans species. Results showed that, urease possessed optimum pH and temperature between 6.5 and 7.0 and 60 and 70° respectively (Krajewska, 2009; Ya Qing FENG, 2005), where the urease relative enzyme activity was higher in germinated beans than the control (mean value).









**Table 1: Result of optimum pH temperature and substrate concentration**

Sample	Germinated			Control		
	pH	T <sup>0</sup>	[S]	pH	T <sup>0</sup>	[S]
<i>Glycine max</i>	6.5	60	0.4	6.5	60	0.4
<i>Mucuna pruriens</i>	7	60	0.4	7	60	0.4
<i>Kerstings geocarpa</i>	6.5	60	0.4	7	70	0.4
<i>Vigna mungo</i>	7	60	0.4	6.5	60	0.4

The effect of urea concentration on the activity of urease from germinated and control of the beans was examined (Figures 1-4). The urease from germinated bean specie exhibited higher activity ( $V_{max}$ ) than the control (Figures 5-8 and Table 2).





However, urease from control showed lower  $K_m$ , when compared to the germinated. This result showed that the control bean is much more sensitive to substrate than the germinated, but with germinated beans, there is higher urea hydrolysis.


**Table 2: Kinetic parameters of urease enzyme from germinated and ungerminated beans**

Sample	$K_m$ (Germinated) mM	$V_{max}$ (Germinated) $\mu\text{mol}/\text{smg}^{-1}$	$K_m$ (Control) mM	$V_{max}$ (Control) $\mu\text{mol}/\text{smg}^{-1}$
<i>Glycine max</i>	0.45	$13.33 \times 10^{-4}$	0.21	$2.00 \times 10^{-4}$
<i>Mucuna pruriens</i>	0.09	$11.34 \times 10^{-4}$	0.05	$11.04 \times 10^{-4}$
<i>Kerstings geocarpa</i>	0.13	$3.04 \times 10^{-4}$	0.1	$1.18 \times 10^{-4}$
<i>Vigna mungo</i>	0.18	$4.90 \times 10^{-4}$	0.02	$0.17 \times 10^{-4}$

The behavior of the reaction velocity ( $V_{max}$ ) and the Michelis-Menten constant ( $K_m$ ) from substrate concentration profile ( $[S]$ ) evaluated. From figures 1-8, it is seen that the reaction velocity was lower in urease from control when compared ( $p < 0.05$ ) to germinated beans at the same substrate concentration. With substrate concentration increasing, the reaction velocity increased, and then kept constant with a maximum value when substrate concentration increased to a certain value (figures showing substrate concentration profile), which fulfilled the basic conditions of Michaelis-Menten equation. Therefore the kinetic parameters are in accordance with Michelis-Menten rate reaction. On a general note, kinetic parameter varies with substrate concentration profile and dilution


factor of the enzyme. Notwithstanding, Michelis-Menten constants for enzymes usually range between  $10^{-2}$  and  $10^{-5}$  mM for most enzymes (Wolfgang, 2007).

Additionally, results (Figures 9 and 10) show that, germination increased the  $K_m$  and  $V_{max}$  of the enzyme when compared ( $p < 0.05$ ) to the control. This feature might be due to the fact that, urease enzyme under germination was metabolically active by making use of seedling's conserved protein to properly fix nickel at the active site of the enzyme to enhance increased hydrolysis of internally generated and external urea (fertilizer) (Polacco & Holland, 1994; Mobley et al., 1995).



Metabolically active urease possessed higher activity ( $V_{max}$ ) in order to facilitate hydrolysis of high concentration of urea in plant under germination and also prevent leaf necrosis which usually occurs owing to accumulation of urea in growing plant (Eskew et al., 1983; Krogmeier et al., 1989; Stebbins et al., 1991). This phenomenon reduced the sensitivity (high  $K_m$ ) of the enzyme for urea compared

( $p < 0.05$ ) to the control. The urease from control beans (non-germinated) was not performing any metabolic activity, it remain inactive in its natural state before it was isolated. As result it happen to be more sensitive because urea concentration was lower than that of the germinated beans.



## Conclusion

The result of this work shows that, germination tend to increase the  $V_{max}$  and  $K_m$  of urease, thereby increasing the activity of the enzyme for higher concentration of urea, most especially in growing plant, and also making the enzyme moderately sensitive to substrate.

## Acknowledgement

I want to express my profound and sincere appreciation to Prof. Gbodi and Mr. Abdujeleel for their technical and moral support given to me while working in the Biochemistry Department Laboratory at Ibrahim Badamoshi Babangida University, Lapai, Niger State, Nigeria.

## Reference

- [1] Arvind M. Kayastha and Nilanjana Das, (1999). *Biochemical education* 27: 114-117.
- [2] Eskew, D.L., R.M. Welch, and E.E. Cary. 1983. Nickel: an essential micronutrient for legumes and possibly all higher plants. *Science* **222**: 621-226.
- [3] Fujinawa, S.B.G and Dela T.P. (1990). Appliatio of acid urease to reduction of urea in commercial wines. *Am J. Enol. Viticul.*, 42: 350-354.
- [4] Fumuyiwa, OO and Ouch, CS (1991). Modification of acid activity by fluoride ions and malic acid in wines. *Am. J. Enol. Viticul.*, 91: 350-359
- [5] Gerenda's J, Sattelmacher B (1999). Influence of Ni supply on growth and nitrogen metabolism of *Brassica napus* L. grown with  $NH_4NO_3$  or urea as N source. *Ann Bot* **83**: 65-71.
- [6] Krogmeier, M.J., G.W. McCarty, and J.M. Bremner. (1989). Phtotoxicity of foliar-applied urea. *Proc. Natl. Acad. Sci. USA* **86**: 9189.
- [7] Mahmood, A. (2001). Investigation of the reversible inhibition of Butrylcholinesterase by mercury chloride. *J. Med. Sci.*, 1, 251-254.
- [8] Moblely, H.L., Island, M.D. & Hausinger, R.P. (1995). Molecular biology of microbial ureases. modified polucarbonate membrane. *Biotechnol Appl Biochem.*, 27: 245-248.
- [9] Polacco, J.C. & Holland, M.A. (1994). Genetic control of plant ureases; in *Genetic Engineering* (Setlow, J.K., ed.) vol. 16, pp. 33-48, Plenum Press, New York.
- [10] Preininger, C. (1999). The enzymatic determination of mercury and copper using acid urease. The effect of buffers. *Mikrochim. Acta.* 130, 209-214.
- [11] Segel, I. H. (1975). *Enzyme Kinetics*. Wiley, New York.
- [12] Stebbins, N., M.A. Holland, S.R. Cianzio, and J.C. Polacco. 1991. Genetics tests of the roles of the embryonic urease of soybean. *Plant Physiol.* **97**: 1004.
- [13] Sumner, J. B. (1926). The isolation and crystallization of the enzyme urease. *J. Biol. Chem.* 69, 435-441.
- [14] Tsai, H.; Doong, R. (2005) Simultaneous determination of pH, urea, acetylcholine and heavy metals using array based enzymatic optical biosensor. *Biosens. Bioelectron.* 20, 1796-1804.
- [15] Whitaker, J.R. and Granum, P.E. (1980). An absolute method for protein determination based on difference of absorbance at 260 and 280nm. *Anal Biochem.* 109: 156-159.
- [16] Wikipedia, (2006). Urea from Wikipedia, the free encyclopedia 14: GNU free documentation license. Wikipedia is a registerd trademark of Wikimedia foundation, inc. version 1.2 november 2002. Copyright (c) 2000, 2001, 2002 free software foundation, USA., Boston, MA 02110-1301.
- [17] Wolfgang A., (2007). *Enzyme in industry* third edition. ISBN: 978-3-527-31689-2.
- [18] Won, K., Kim, S., Kim, K.J., Park, H.W and Moon, S.J. (2005) *Process Biochem.* **40**: 2149.
- [19] Y. Kulshrestha, Q. Husain, (2006). *Enzyme Microb. Technol.* **38**: 47.
- [20] Ya Qing FENG (2005). *Chinese Chemical Letters* Vol. 16, No. 1, pp 135-138.  
<http://www.imm.ac.cn/journal/ccl.html>
- [21] Zhu, Z. L., Chen, D, L. (2002). Nitrogen fertilizer use in China-Contributions to food production, impact on environment and best management strategies. *Nutri Cycl Agroecosys* 63, 117-127.